

AD-A210 132

THE FILE COPY

②

OFFICE OF NAVAL RESEARCH

Contract N00014-86-K-0043

TECHNICAL REPORT No. 104

Squeezing of Atomic Variables in the One-Photon and Two-Photon  
Jaynes-Cummings Model

by

Xiao-shen Li, D. L. Lin and Thomas F. George

Prepared for Publication

in

Physical Review A

Departments of Chemistry and Physics  
State University of New York at Buffalo  
Buffalo, New York 14260

June 1989

Reproduction in whole or in part is permitted for any purpose of the  
United States Government.

This document has been approved for public release and sale;  
its distribution is unlimited.

DTIC  
ELECTE  
JUL 17 1989  
S B D  
Cb

89

7

17

102

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

## REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UBUFFALO/DC/89/TR-104			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Depts. Chemistry & Physics State University of New York		6b. OFFICE SYMBOL (If applicable)		7a. NAME OF MONITORING ORGANIZATION	
6c. ADDRESS (City, State, and ZIP Code) Fronczak Hall, Amherst Campus Buffalo, New York 14260		7b. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217			
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Contract N00014-86-K-0043	
8c. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217		10. SOURCE OF FUNDING NUMBERS			
		PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Squeezing of Atomic Variables in the One-Photon and Two-Photon Jaynes-Cummings Model					
12. PERSONAL AUTHOR(S) Xiao-shen Li, D. L. Lin and Thomas F. George					
13a. TYPE OF REPORT		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) June 1989	
				15. PAGE COUNT 17	
16. SUPPLEMENTARY NOTATION Prepared for publication in Physical Review A					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	JAYNES-CUMMINGS MODEL		
			ATOMIC DIPOLE MOMENT		
			SQUEEZING		
			> ONE-PHOTON PROCESS, TWO-PHOTON PROCESS, QUANTUM MECHANICAL STUDY		
19. ABSTRACT (Continue on reverse if necessary and identify by block number)  Squeezing of the atomic dipole moment in the one-photon and two-photon Jaynes-Cummings model is investigated with different initial conditions.  Almost perfect squeezing is found in certain cases.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson			22b. TELEPHONE (Include Area Code) (202) 696-4410		22c. OFFICE SYMBOL

Physical Review A, in press

Squeezing of atomic variables in the one-photon and two-photon  
Jaynes-Cummings model

Xiao-shen Li  
Center for Theoretical Physics  
Chinese Center of Advanced Science and Technology (World Laboratory)  
P. O. Box 8730  
Beijing 100080, People's Republic of China

and

Shanghai Institute of Metallurgy  
Chinese Academy of Sciences  
Shanghai, People's Republic of China

D. L. Lin and Thomas F. George  
Department of Physics & Astronomy  
239 Fronczak Hall  
State University of New York at Buffalo  
Buffalo, New York 14260

Abstract

Squeezing of the atomic dipole moment in the one-photon and two-photon  
Jaynes-Cummings model is investigated with different initial conditions.  
Almost perfect squeezing is found in certain cases.

PACS Numbers: 32.80.-t, 42.5.Kb

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

## I. Introduction

It is by now well known that squeezed light has potential applications in low-noise communications and high-precision measurements involving light.<sup>1,2</sup> The Jaynes-Cummings (JC) model<sup>3,4</sup> is one of the possible nonlinear optical models capable of creating squeezed states.<sup>5</sup> A number of calculations of field squeezing have been made in recent years. In the standard JC model, the maximum field squeezing was found to be 20%,<sup>5</sup> but in a recent paper it was argued that this squeezing could be 100%.<sup>6</sup> When the initial coherent state of the field is replaced by the vacuum state, the maximum squeezing is increased to 25%.<sup>7</sup> More squeezing is found in many other cases of the generalized JC model. The maximum squeezing is 42% for two-level two-photon,<sup>8</sup> 57% for two-level three-photon,<sup>9</sup> 31% for three-level one-mode,<sup>10</sup> 36% for four-level one-mode,<sup>11</sup> and 52% for ten-level one-mode<sup>12</sup> cases. However, little attention has been paid to the squeezing of the atomic variables. The squeezing of the atomic dipole operators for a two-level atom driven by a classical field<sup>13</sup> has been investigated. A general relation has been established between the coherent states of the SU(2) and SU(1,1) Lie algebra and possible reduction of fluctuations in angular momentum.<sup>14</sup> For the special case of a vacuum initial field mode, the relation between the atomic variable squeezing and the field squeezing has been examined.<sup>15</sup> Only recently, a quantum mechanical study devoted to the squeezing of the atomic dipole moment has been carried out,<sup>16</sup> where it is found that a maximum squeezing of 60% can be reached.

In this note we calculate the squeezing of the atomic dipole moment in the one-photon and two-photon JC model with different initial conditions. It is found that the atomic variables generally possess stronger squeezing than the field under the same conditions. In particular circumstances, a nearly pure squeezing state can be achieved. In the standard JC model, a one-mode

cavity field with frequency  $\Omega$  couples to a two-level atom through the well-known Hamiltonian in the rotating-wave approximation,

$$H = \frac{1}{2} \hbar \omega \sigma_3 + \hbar \Omega a^\dagger a + \hbar \lambda (a^\dagger \sigma^- + a \sigma^+) \quad , \quad (1)$$

where  $a^\dagger(a)$  and  $\Omega$  are the creation (annihilation) operator and frequency of the cavity field, respectively, and  $\lambda$  is the coupling constant. The two-level atom with transition frequency  $\omega$  is described by the Pauli raising and lowering operators  $\sigma^\pm$  and the inversion operator  $\sigma_3 = \sigma^+ \sigma^- - \sigma^- \sigma^+$ . The dispersive and absorptive components of the slowly-varying atomic dipole<sup>17</sup> can be written as

$$\sigma_1 = \frac{1}{2} (\sigma^+ e^{-i\omega t} + \sigma^- e^{i\omega t}) \quad (2)$$

and

$$\sigma_2 = \frac{1}{2i} (\sigma^+ e^{-i\omega t} - \sigma^- e^{i\omega t}) \quad , \quad (3)$$

respectively. They obey the commutation relation

$$[\sigma_1, \sigma_2] = \frac{i}{2} \sigma_3 \quad (4)$$

and the corresponding uncertainty relation

$$(\Delta \sigma_1)^2 (\Delta \sigma_2)^2 \geq \frac{1}{16} \langle \sigma_3 \rangle^2 \quad . \quad (5)$$

The atomic state is said to be squeezed when  $\sigma_1$  or  $\sigma_2$  satisfies the relation

$$(\Delta\sigma_i)^2 < \frac{1}{4} |\langle\sigma_3\rangle|, \quad i = 1, 2. \quad (6)$$

Since

$$(\Delta\sigma_1)^2 = \frac{1}{4} - (\text{Re}\langle\sigma^-\rangle e^{i\omega t})^2 \quad (7a)$$

$$(\Delta\sigma_2)^2 = \frac{1}{4} - (\text{Im}\langle\sigma^-\rangle e^{i\omega t})^2, \quad (7b)$$

the condition described by Eq. (6) can be rewritten as

$$S_1 = \frac{1 - 4(\text{Re}\langle\sigma^-\rangle e^{i\omega t})^2}{|\langle\sigma_3\rangle|} < 1 \quad (8a)$$

or

$$S_2 = \frac{1 - 4(\text{Im}\langle\sigma^-\rangle e^{i\omega t})^2}{|\langle\sigma_3\rangle|} < 1 \quad (8b)$$

for squeezing in the dispersive or absorptive component of the dipole. In what follows we shall investigate this squeezing effect by using different initial conditions for the atom and the cavity field.

## II. Squeezing in the JC model

We denote by  $|+\rangle$  and  $|-\rangle$  the excited and ground states of the atom and by  $|n\rangle$  the Fock states of the field. First we consider the initial condition

that the atom is in the excited state and the field in a two-photon coherent state or squeezed state,<sup>1,2,18</sup>

$$|\alpha, r\rangle = \sum_n F(n) |n\rangle \quad (9)$$

$$F(n) = (n! \cosh r)^{-1/2} \left( \frac{\beta}{2 \cosh r} \right)^n Y^{-n} H_n(Y) \exp\left[-\frac{\beta^2}{2}(1 - \tanh r)\right] \quad (10)$$

where  $\beta = \alpha e^r$  and  $Y = \beta(2 \cosh r \sinh r)^{-1/2}$ , and we have chosen the squeezing parameter  $r$  to be real.<sup>19</sup> The initial mean photon number is given by  $\bar{n} = |\alpha|^2 + \sinh^2 r$ , so that the initial state vector for the atomic-field system can be written as

$$|\psi(0)\rangle = \sum_n F(n) |+, n\rangle \quad (11)$$

At a time  $t > 0$ , the state vector in the interaction picture is found from the JC model Hamiltonian (1) to be

$$|\psi(t)\rangle = \sum_n F(n) [A(n, t) |+, n\rangle + B(n+1, t) |-, n+1\rangle] \quad (12)$$

where

$$A(n, t) = e^{-i\Delta t/2} (\cos \mu t + i \frac{\Delta}{2} \sin \mu t / \mu) \quad (13)$$

$$B(n+1, t) = -i \frac{Y}{\mu} e^{-i\Delta t/2} \sin \mu t \quad (14)$$

with  $V = \lambda\sqrt{n+1}$ ,  $\mu^2 = V^2 + \Delta^2/4$  and  $\Delta = \Omega - \omega$ . From Eqs. (12)-(14), we obtain the expectation values of the atomic operators:

$$\begin{aligned} \langle \sigma^- \rangle e^{i\omega t} &= \langle \psi(t) | \sigma^- e^{i\omega t} | \psi(t) \rangle \\ &= \sum_{n=0}^{\infty} F^*(n) F(n+1) B^*(n+1) A(n+1) \end{aligned} \quad (15)$$

$$\langle \sigma_3 \rangle = \sum_{n=0}^{\infty} |F(n)|^2 [ |A(n)|^2 - |B(n+1)|^2 ] \quad (16)$$

The time evolutions of  $S_1$  and  $S_2$  are calculated for the case of off-resonant excitation with medium strength. The units employed throughout this paper are  $\lambda$  for energy and  $\lambda^{-1}$  for time. The results are presented in Fig. 1. It is observed that  $\sigma_1$  and  $\sigma_2$  are squeezed alternatively. Roughly speaking, the squeezing parameters  $S_1$  and  $S_2$  oscillate at about the same frequency but out of phase. A more careful analysis reveals that both are composed of two oscillations with different frequencies. The maximum squeezing in this case is approximately 25% for both  $S_1$  and  $S_2$ .

For the case of on-resonance excitation,  $\langle \sigma_1 \rangle$  is zero but its fluctuation is still nonvanishing because of quantum mechanical effects. Figure 2 shows the time evolution of  $S_2$  for the on-resonance excitation by a squeezed and a coherent light field. A comparison of (a) and (b) in Fig. 2 shows that squeezing occurs only for a short time after the interaction is turned on when the initial field is in a coherent state. The maximum squeezings in these cases are, however, about the same, namely, 65% and 68% for the coherent and squeezed excitations, respectively. The role played by



the detuning can be clearly seen by comparing Figs. 1(b) and 2(a). Evidently, a larger detuning implies weaker coupling between the atom and the field and hence less squeezing. Although the results shown are obtained for the same  $\bar{n}$ , we have found from our numerical study that in general a larger exciting strength or larger  $\bar{n}$  results in stronger squeezing effects.

Next we consider a vacuum initial field in the JC model. The atom is injected into the field in a coherent superposition of excited and ground states,<sup>7</sup>

$$|\psi(0)\rangle = e^{i\phi} \cos\theta |-,0\rangle + \sin\theta |+,0\rangle . \quad (17)$$

From the Hamiltonian (1) and Eq. (17), the state vector at time  $t > 0$  can be written as

$$|\psi(t)\rangle = C_0^- |-,0\rangle + C_1^- |-,1\rangle + C_0^+ |+,0\rangle , \quad (18)$$

where

$$C_1^-(t) = -i \sin\theta \sin\lambda t \quad (19)$$

$$C_0^+(t) = \sin\theta \cos\lambda t \quad (20)$$

$$C_0^-(t) = e^{i\phi} \cos\theta . \quad (21)$$

Using the same procedure as outlined above, we can write

$$\langle \sigma^- \rangle e^{i\omega t} = \langle \psi(t) | \sigma^- e^{i\omega t} | \psi(t) \rangle = C_0^{-*} C_0^+ - \frac{1}{2} e^{-i\phi} \sin 2\theta \cos \lambda t \quad (22)$$

and

$$\langle \sigma_3 \rangle = 1 - 2\sin^2\theta \cos^2\lambda t \quad . \quad (23)$$

Inserting Eqs. (22) and (23) into (8), we find the squeezing parameters as

$$S_1 = (1 - \sin^2 2\theta \cos^2 \phi \cos^2 \lambda t) / |1 - 2\sin^2 \theta \cos^2 \lambda t| \quad , \quad (24)$$

$$S_2 = (1 - \sin^2 2\theta \sin^2 \phi \cos^2 \lambda t) / |1 - 2\sin^2 \theta \cos^2 \lambda t| \quad . \quad (25)$$

From these two equations one can see that when the atom starts in a purely excited state, i.e.,  $\theta = \frac{\pi}{2}$ , there is no squeezing effect. It is apparent that maximum squeezing occurs when  $\lambda t = k\pi$  ( $k = 0, 1, 2, \dots$ ) and  $\phi = k\pi$  (for  $\sigma_1$ ) or  $(k + \frac{1}{2})\pi$  (for  $\sigma_2$ ). We have, under these conditions,

$$S_1 = (1 - \sin^2 2\theta) / |1 - 2\sin^2 \theta| = |\cos 2\theta| \quad , \quad (26)$$

where  $\theta = \frac{1}{2}(k + \frac{1}{2})\pi$ . Equation (20) yields

$$S_1 = 0 \quad . \quad (27)$$

This means that the maximum squeezing of 100% is possible.

### III. Squeezing in two-photon processes

A simple model for atom-radiation interaction that can be solved analytically has recently been proposed.<sup>7</sup> It involves two-photon Raman

coupling between two atomic states  $|+\rangle$  and  $|-\rangle$  degenerate in energy. The Hamiltonian for this model is therefore

$$H = \hbar\Omega a^\dagger a + \hbar\lambda a^\dagger a(\sigma^- + \sigma^+) \quad , \quad (28)$$

where we have set the atomic transition frequency to be zero because of the degeneracy. It has been shown that the system has perfect revival.<sup>7</sup> Here we shall study the squeezing of the atomic dipole operator in this model.

We assume that the atom starts in the initial state  $|+\rangle$  and the field is in a coherent state  $|\alpha\rangle$ ,

$$|\alpha\rangle = \sum_n G(n) |n\rangle \quad (29)$$

$$G(n) = \frac{\alpha^n}{\sqrt{n!}} e^{-\bar{n}/2} \quad , \quad (30)$$

where  $\bar{n} = |\alpha|^2$  is the initial mean photon number of the field. The initial state vector then takes the form

$$|\psi(0)\rangle = \sum_n G(n) |+, n\rangle \quad . \quad (31)$$

At any time  $t > 0$ , the state vector can be written as

$$|\psi(t)\rangle = \sum_{n=0}^{\infty} G(n) [D^+(n) |+, n\rangle + D^-(n) |-, n\rangle] \quad , \quad (32)$$

where

$$D^+(n,t) = \cos(\lambda nt) \quad (33a)$$

$$D^-(n,t) = -i \sin(\lambda nt) \quad (33b)$$

From Eqs. (32) and (33), one can easily find the mean value of the atomic dipole moment as

$$\langle \sigma^- \rangle = \frac{i}{2} \sum_{n=0}^{\infty} e^{-\bar{n}} \frac{\bar{n}^n}{n!} \sin(2\lambda nt) = \frac{i}{2} \exp[-2\bar{n} \sin^2(\lambda t)] \sin[\bar{n}(\sin 2\lambda t)] \quad (34)$$

and the inversion

$$\langle \sigma_3 \rangle = \exp(-2\bar{n} \sin^2 \lambda t) \cos(\bar{n} \sin 2\lambda t) \quad (35)$$

It is clear that the mean value of the dispersive component for the atomic dipole fluctuates around zero. Hence there is no squeezing in this component. As for the squeezing in the other component  $\sigma_2$ , we find from Eqs. (8), (34) and (35)

$$S_2 = \frac{1 - \exp(-4\bar{n} \sin^2 \lambda t) \sin^2(\bar{n} \sin 2\lambda t)}{\exp(-2\bar{n} \sin^2 \lambda t) |\cos(\bar{n} \sin 2\lambda t)|} \quad (36)$$

To find the maximum magnitude of squeezing, we may consider the case where  $\bar{n} \gg 1$  but  $\bar{n}(\lambda t)^2 \ll 1$ . Thus we have

$$S_2 \approx \frac{1 - [1 - 4\bar{n}(\lambda t)^2] \sin^2(2\bar{n}\lambda t)}{|\cos(2\bar{n}\lambda t)|}$$

$$= \frac{1 - (1 - \bar{n}^{-1}x) \sin^2 x}{|\cos x|}, \quad (37)$$

where  $x = 2\bar{n}\lambda t$ . When  $x$  is in the vicinity of  $\frac{\pi}{2}$ , i.e.,  $x = \frac{\pi}{2} - \epsilon$  with  $\frac{1}{\bar{n}} \ll \epsilon \ll 1$ , we may expand  $\sin x$  and  $\cos x$  in a Taylor series around  $\frac{\pi}{2}$  to the second term. Hence, for  $\lambda t \approx \frac{\pi}{4} \bar{n}^{-1}$  we have

$$S_2 \approx \frac{\pi^2}{4} (\bar{n}\epsilon)^{-1}. \quad (38)$$

Equation (38) implies that for strong excitation,  $S_2$  is very small and the maximum magnitude of squeezing is nearly 100%.

#### Acknowledgments

This research was supported in part by the National Science Foundation under Grant CHE-8620274, the Office of Naval Research and the Air Force Office of Scientific Research (AFSC), United States Air Force, under Contract F49620-86-C-0009. The United States Government is authorized to reproduce and distribute reprints for governmental purposes notwithstanding any copyright notation hereon.

## References

1. H. P. Yuen, Phys. Rev. A 13, 2226 (1976).
2. D. F. Walls, Nature 306, 141 (1983).
3. E. T. Jaynes and F. W. Cummings, Proc. IEEE 51, 89 (1963).
4. J. H. Eberly, N. B. Narozhny, J. J. Sanchez-Mondragon, Phys. Rev. Lett. 44, 1323 (1980).
5. P. Meystre and M. S. Zubairy, Phys. Lett. 89A, 390 (1982).
6. J. R. Kukliński and J. L. Madajczyk, Phys. Rev. A 37, 3175 (1988).
7. P. L. Knight, Phys. Scr. T12, 51 (1986).
8. G. Compagno, J. S. Peng and F. Persico, Opt. Commun. 57, 415 (1986).
9. A. S. Shumovsky, F. Le Kien and E. I. Abskenderov, Phys. Lett. 124A, 351 (1987).
10. S. Y. Zhu, Z. D. Liu and X. S. Li, Phys. Lett. 128A, 89 (1988).
11. Z. D. Liu, S. Y. Zhu and X. S. Li, J. Mod. Opt., in press.
12. F. L. Li, X. S. Li and D. L. Lin, unpublished.
13. D. F. Walls and P. Zoller, Phys. Rev. Lett. 47, 709 (1981).
14. K. Wodkiewicz and J. H. Eberly, J. Opt. Soc. Am. B 2, 458 (1985).
15. K. Wódkiewicz, P. L. Knight, S. J. Buckle and S. M. Barnett, Phys. Rev. A 35, 2567 (1987).
16. X. S. Li, D. L. Lin, T. F. George and Z. D. Liu, Phys. Rev. A, in press.
17. N. B. Narozhny, J. J. Sanchez-Mondragon and J. H. Eberly, Phys. Rev. A 23, 236 (1981).
18. R. Loudon and P. L. Knight, J. Mod. Opt. 34, 709 (1987).
19. G. J. Milburn and D. F. Walls, Phys. Rev. A 27, 392 (1983).

### Figure Captions

1. Time evolution of the squeezing parameters (a)  $S_1$  and (b)  $S_2$  calculated for the case of off-resonance excitation. The parameters used are  $\bar{n} = 100$ ,  $r = 1$  and  $\Delta = 50$ .
2. Time evolution of  $S_2$  for the case of on-resonance excitation by a (a) squeezed ( $r = 1$ ) and (b) coherent ( $r = 0$ ) field with  $\bar{n} = 100$ .







Fig. 1(a)





(b)

Fig. 2



TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
		Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. J. E. Jensen  
Hughes Research Laboratory  
3011 Malibu Canyon Road  
Malibu, California 90265

Dr. C. B. Harris  
Department of Chemistry  
University of California  
Berkeley, California 94720

Dr. J. H. Weaver  
Department of Chemical Engineering  
and Materials Science  
University of Minnesota  
Minneapolis, Minnesota 55455

Dr. F. Kutzler  
Department of Chemistry  
Box 5055  
Tennessee Technological University  
Cookeville, Tennessee 38501

Dr. A. Reisman  
Microelectronics Center of North Carolina  
Research Triangle Park, North Carolina  
27709

Dr. D. DiLella  
Chemistry Department  
George Washington University  
Washington D.C. 20052

Dr. M. Grunze  
Laboratory for Surface Science and  
Technology  
University of Maine  
Orono, Maine 04469

Dr. R. Reeves  
Chemistry Department  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. J. Butler  
Naval Research Laboratory  
Code 6115  
Washington D.C. 20375-5000

Dr. Steven M. George  
Stanford University  
Department of Chemistry  
Stanford, CA 94305

Dr. L. Interante  
Chemistry Department  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. Mark Johnson  
Yale University  
Department of Chemistry  
New Haven, CT 06511-8118

Dr. Irvin Heard  
Chemistry and Physics Department  
Lincoln University  
Lincoln University, Pennsylvania 19352

Dr. W. Knauer  
Hughes Research Laboratory  
3011 Malibu Canyon Road  
Malibu, California 90265

Dr. K.J. Klaubunde  
Department of Chemistry  
Kansas State University  
Manhattan, Kansas 66506

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. G. A. Somorjai  
Department of Chemistry  
University of California  
Berkeley, California 94720

Dr. J. Murday  
Naval Research Laboratory  
Code 6170  
Washington, D.C. 20375-5000

Dr. J. B. Hudson  
Materials Division  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. Theodore E. Madey  
Surface Chemistry Section  
Department of Commerce  
National Bureau of Standards  
Washington, D.C. 20234

Dr. J. E. Demuth  
IBM Corporation  
Thomas J. Watson Research Center  
P.O. Box 218  
Yorktown Heights, New York 10598

Dr. M. G. Lagally  
Department of Metallurgical  
and Mining Engineering  
University of Wisconsin  
Madison, Wisconsin 53706

Dr. R. P. Van Duyne  
Chemistry Department  
Northwestern University  
Evanston, Illinois 60637

Dr. J. M. White  
Department of Chemistry  
University of Texas  
Austin, Texas 78712

Dr. D. E. Harrison  
Department of Physics  
Naval Postgraduate School  
Monterey, California 93940

Dr. R. L. Park  
Director, Center of Materials  
Research  
University of Maryland  
College Park, Maryland 20742

Dr. W. T. Peria  
Electrical Engineering Department  
University of Minnesota  
Minneapolis, Minnesota 55455

Dr. Keith H. Johnson  
Department of Metallurgy and  
Materials Science  
Massachusetts Institute of Technology  
Cambridge, Massachusetts 02139

Dr. S. Sibener  
Department of Chemistry  
James Franck Institute  
5640 Ellis Avenue  
Chicago, Illinois 60637

Dr. Arnold Green  
Quantum Surface Dynamics Branch  
Code 3817  
Naval Weapons Center  
China Lake, California 93555

Dr. A. Wold  
Department of Chemistry  
Brown University  
Providence, Rhode Island 02912

Dr. S. L. Bernasek  
Department of Chemistry  
Princeton University  
Princeton, New Jersey 08544

Dr. W. Kohn  
Department of Physics  
University of California, San Diego  
La Jolla, California 92037

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. F. Carter  
Code 6170  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. Richard Colton  
Code 6170  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. Dan Pierce  
National Bureau of Standards  
Optical Physics Division  
Washington, D.C. 20234

Dr. R. Stanley Williams  
Department of Chemistry  
University of California  
Los Angeles, California 90024

Dr. R. P. Messmer  
Materials Characterization Lab.  
General Electric Company  
Schenectady, New York 22217

Dr. Robert Gomer  
Department of Chemistry  
James Franck Institute  
5640 Ellis Avenue  
Chicago, Illinois 60637

Dr. Ronald Lee  
R301  
Naval Surface Weapons Center  
White Oak  
Silver Spring, Maryland 20910

Dr. Paul Schoen  
Code 6190  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. John T. Yates  
Department of Chemistry  
University of Pittsburgh  
Pittsburgh, Pennsylvania 15260

Dr. Richard Greene  
Code 5230  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. L. Kesmodel  
Department of Physics  
Indiana University  
Bloomington, Indiana 47403

Dr. K. C. Janda  
University of Pittsburgh  
Chemistry Building  
Pittsburg, PA 15260

Dr. E. A. Irene  
Department of Chemistry  
University of North Carolina  
Chapel Hill, North Carolina 27514

Dr. Adam Heller  
Bell Laboratories  
Murray Hill, New Jersey 07974

Dr. Martin Fleischmann  
Department of Chemistry  
University of Southampton  
Southampton SO9 5NH  
UNITED KINGDOM

Dr. H. Tachikawa  
Chemistry Department  
Jackson State University  
Jackson, Mississippi 39217

Dr. John W. Wilkins  
Cornell University  
Laboratory of Atomic and  
Solid State Physics  
Ithaca, New York 14853

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. R. G. Wallis  
Department of Physics  
University of California  
Irvine, California 92664

Dr. D. Ramaker  
Chemistry Department  
George Washington University  
Washington, D.C. 20052

Dr. J. C. Hemminger  
Chemistry Department  
University of California  
Irvine, California 92717

Dr. T. F. George  
Chemistry Department  
University of Rochester  
Rochester, New York 14627

Dr. G. Rubloff  
IBM  
Thomas J. Watson Research Center  
P.O. Box 218  
Yorktown Heights, New York 10598

Dr. Horia Metiu  
Chemistry Department  
University of California  
Santa Barbara, California 93106

Dr. W. Goddard  
Department of Chemistry and Chemical  
Engineering  
California Institute of Technology  
Pasadena, California 91125

Dr. P. Hansma  
Department of Physics  
University of California  
Santa Barbara, California 93106

Dr. J. Baldeschwieler  
Department of Chemistry and  
Chemical Engineering  
California Institute of Technology  
Pasadena, California 91125

Dr. J. T. Keiser  
Department of Chemistry  
University of Richmond  
Richmond, Virginia 23173

Dr. R. W. Plummer  
Department of Physics  
University of Pennsylvania  
Philadelphia, Pennsylvania 19104

Dr. E. Yeager  
Department of Chemistry  
Case Western Reserve University  
Cleveland, Ohio 41106

Dr. N. Winograd  
Department of Chemistry  
Pennsylvania State University  
University Park, Pennsylvania 16802

Dr. Roald Hoffmann  
Department of Chemistry  
Cornell University  
Ithaca, New York 14853

Dr. A. Steckl  
Department of Electrical and  
Systems Engineering  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. G.H. Morrison  
Department of Chemistry  
Cornell University  
Ithaca, New York 14853